

BE PREPARED.

LEARN ABOUT VACCINE PREVENTION

COLLEGE IMMUNIZATIONS

In compliance with Oregon Law Senate Bill 274, we are providing you with required information on vaccine preventable diseases known to occur in individuals 16-21 years of age.

Make sure you are up-to-date on all your vaccinations. Any student entering a public or private postsecondary education institution in this state for the first time after July 1, 2008, shall, within forty-five days after the start of classes, present to the appropriate institution certification from a licensed physician that the student has received, or is in the process of receiving, the required two doses of immunization against measles, mumps, and rubella. Additional vaccinations (or documentation of these vaccinations or laboratory proof of immunity) may be required for student admission to certain health profession programs. Immunizations for tetanus, diphtheria, poliomyelitis, hepatitis B, varicella and meningitis are recommended, as is a tuberculin test. An annual influenza vaccination is recommended, particularly for students living in residence halls, to minimize disruption of routine activities during influenza out- breaks.

REQUIRED:

- Measles (usually obtained in the form MMR)

RECOMMENDED:

- Meningococcal B
- Meningococcal ACWY
- Tdap *
- Varicella *
- Hepatitis B *
- HPV
- Influenza

*certain vaccines may be required for students in the health profession programs

Why get vaccinated?

MMR

Measles, mumps, and rubella are serious diseases. Before vaccines they were very common, especially among children.

MEASLES

Signs & Symptoms

Rash, cough, runny nose, eye irritation, fever, ear infection, pneumonia, seizures (jerking and staring), brain damage, and death.

MUMPS

Signs & Symptoms

Fever, headache, muscle pain, loss of appetite, and swollen glands, deafness, meningitis (infection of the brain and spinal cord covering), painful swelling of the testicles or ovaries, and rarely sterility.

RUBELLA (German Measles)

Signs & Symptoms

Rash, arthritis (mostly in women), mild fever, miscarriage

Measles, mumps, and rubella (MMR) vaccine can protect children (and adults) from all three of these diseases.

HEPATITIS B

Hepatitis B is a serious disease that affects the liver. It is caused by the hepatitis B virus. Hepatitis B can cause mild illness lasting a few weeks, or it can lead to a serious, lifelong illness.

Signs & Symptoms

Fever, fatigue, loss of appetite, nausea, and/or vomiting, jaundice (yellow skin or eyes, dark urine, clay-colored bowel movements), pain in muscles, joints, and stomach, liver damage (cirrhosis), liver cancer, death

Hepatitis B vaccine can prevent hepatitis B and its consequences, including liver cancer and cirrhosis.

MENINGOCOCCAL

Meningococcal disease often occurs without warning — even among people who are otherwise healthy.

Anyone can get meningococcal disease but certain people are at increased risk, including:

Infants younger than one year old, adolescents and young adults 16 through 23 years old, people with certain medical conditions

that affect the immune system, people at risk because of an outbreak in their community.

Signs & Symptoms

Fever, headache, and stiff neck, nausea, vomiting, photophobia (increased sensitivity to light), altered mental status (confusion).

Tdap

Tetanus, diphtheria and pertussis are very serious diseases. Tdap vaccine can provide protection against these disease, and Tdap vaccine given to pregnant women can protect newborn babies against pertussis.

TETANUS (Lockjaw)

Signs & Symptoms

•painful muscle tightening and stiffness, usually all over the body, tightening of muscles in the head and neck blocking the ability to open the mouth, swallow, or sometimes even breathe.

DIPHTHERIA

Signs & Symptoms

Can cause a thick coating to form in the back of the throat, breathing problems, heart failure, paralysis, and death.

PERTUSSIS

Signs & Symptoms

Pertussis (Whooping Cough) causes severe coughing spells, which can cause difficulty breathing, vomiting and disturbed sleep.

Weight loss, incontinence, and rib fractures, pneumonia or death

VARICELLA

Varicella (also called chickenpox) is a common childhood disease. It is usually mild, but it can be serious, especially in young infants and adults.

Signs & Symptoms

Rash, itching, fever, and tiredness, severe skin infection, scars, pneumonia, brain damage, or death.

A person who has had chickenpox can get a painful rash called shingles years later.

Chickenpox vaccine can prevent chickenpox. Most people who get chickenpox vaccine will not get chickenpox. But if someone who has been vaccinated does get chickenpox, it is usually very mild. They will have fewer blisters, are less likely to have a fever, and will recover faster.

HPV

HPV vaccine prevents infection with human papillomavirus (HPV) types that are associated with many cancers, including: cervical cancer in females, vaginal and vulvar cancers in females, anal cancer in females and males, throat cancer in females and males, and penile cancer in males. In addition, HPV vaccine prevents infection with HPV types that cause genital warts in both females and males.

**Vaccination is not a substitute for cervical cancer screening. This vaccine does not protect against all HPV types that can cause cervical cancer. Women should still get regular Pap tests.*

INFLUENZA

Influenza (also called Respiratory Flu) is caused by influenza viruses, and is spread mainly by coughing, sneezing, and close contact. Anyone can get flu. Flu strikes suddenly and can last several days.

Symptoms vary by age, but can include:

Fever/chills, sore throat, muscle aches, fatigue, cough, headache, runny or stuffy nose, pneumonia and blood infections, and cause diarrhea and seizures in children. Medical conditions, such as heart or lung disease, can be worsened by the flu.

Flu is more dangerous for some people. Infants and young children, people 65 years of age and older, pregnant women, and people with certain health conditions or a weakened immune system are at greatest risk.

Flu vaccine can prevent the flu or make it less severe, and prevent the spread of flu to others.